

Product information

【Drug Facts】



〈Kyowa-Gifu-Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs

Rhinitis medication

(TABLETS)

NOSUPOL BIEN CHEWABLE JYOU II

”NOSUPOL Bein Chewable Jyou II” is a well-balanced combination of five active ingredients that effectively improves sneezing, runny nose, nasal congestion, etc.

<i>Active ingredients (in 6 tablets/daily dose)</i>	<i>Purpose</i>
Phenylephrine hydrochloride 30mg*****	Treatment for stuffy nose
Chlorpheniramine maleate 9mg*****	Antihistamine
Belladonna Total Alkaloid 0.45mg*****	Nasal mucus secretion regulation
Glycyrrhizinic acid 30mg*****	Soothe sore throat and cough
Anhydrous Caffeine 60mg*****	Sleepiness prevention

Uses

Alleviation of the following symptoms by acute rhinitis, allergic rhinitis or sinusitis: sneezing, runny nose (too much nasal discharge), stuffy nose, watery eyes, sore throat, headache (heavy head).

Warnings

Do not use for children under 7 years old.

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

- 1. While taking this medicine, please do not take any of the following medicines.**

Other internal medicines for rhinitis or containing antihistamines (cold medicine, antitussives, motion sickness medicine, medicine for allergy, etc.), gastrointestinal analgesic spasm.

- 2. Do not drive a motor vehicle or operate machinery after using this medicine. (Symptoms such as drowsiness, blurred vision, dazzling sunlight may occur.)**
- 3. Do not take long-term continuously.**

[Consultation]

1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A pregnant woman or a woman suspected of being pregnant.
- (3) Senior persons.
- (4) A person who have ever had allergic symptoms due to any medication.
- (5) A person with the following symptoms:
High fever、 Dysuria.
- (6) A person diagnosed as having the following:
Glaucoma, Diabetes, Thyroid dysfunction, Heart disease, Hypertension.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Skin	Rash · Redness、 Itching
Digestive organ	Nausea · Vomiting, Loss of Appetite
Central nervous system	Headache
Urinary system	Dysuria
Others	Face hot flashes, Abnormal dazzling

In very rare cases, the following serious symptoms may occur. In this case, please get medical attention immediately.

Symptoms' name	Symptoms
Aplastic anemia	Bruising, nosebleeds, gum bleeding, fever, skin and mucous membranes appear pale, fatigue, palpitations, shortness of breath, mood worsening, bloodsheds etc. appear.
Agranulocytosis	Sudden high fever, chilling, sore throat etc. appear.

3. The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Thirst, sleepiness, constipation and blurred eyes.

4. If these symptoms do not improve after having used this medicine 5 or 6 times, stop taking the medicine, and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Directions

Dissolve or chew the following daily dose on three times a day in the mouth. However, please take at least 4 hours between doses.

Age	One dose	Number of doses per day
adult and children 15 years and over	2 tablets	3 times
children 7 years to under 15 years	1 tablet	
children under 7 years	Do not use	

[When using this product]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) When using this medicine for the first time.

Please push the projection of the PTP sheet containing tablets strongly with your fingertips as shown below, break the aluminum sheet on the back side, push out and take it.

(If you mistakenly swallow the PTP sheet as it is, it leads to an unexpected accident such as stabbing into the esophagus mucosa)



- (4) Take care not to break the tablets when taking out of the PTP sheet.
- (5) Do not handle this medicine with wet hands.

Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)

(4) Please do not take products that have passed the expiration date

Inactive ingredients:

Lactose, Sucralose, Polyvinyl alcohol,
Acrylic acid, Methyl methacrylate copolymer,
Xylitol, Croscopidone, Al silicate, Cellulose,
Silicon dioxide, Magnesium stearate,
Thaumatococcus, D-Mannitol, Hydrogen phosphate
Ca, Contains hydroxypropyl starch,
Peppermint oil and Cross-CMC-Na.

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.