

Product information

【Drug Facts】



<Kyowa-Gifu-Japan>

For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Class 2 drugs **Medicine for motion sickness (LIQUID)**

NEW ARTAR LIQUID

- "NEW ARTAR LIQUID", taken once a day, reduce nervous excitement caused by vehicle sway and acceleration, and reduce or relieve the occurrence of dizziness, nausea and headache caused by motion sickness. If you take about 30 minutes before getting on a car, bus, ship, airplane etc, you will be relieved from unpleasant motion sickness, thereby making travel and hobbies more fun.

<i>Active ingredient (in 2 bottles/daily dose)</i>	<i>Purpose</i>
(in 40mL/2 bottles)	
Scopolamine Hydrobromide Hydrate 0.48mg	Sedative
Chlorpheniramine maleate 8mg.....	Anti-allergy
Anhydrous Caffeine 100mg	Headache treatment

Uses For the prevention and treatment of dizziness, nausea, headache associated with motion sickness.

Warnings Do not use for children.

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

1 While taking this medicine, please do not take any of the following medicines.

Other medicines such as motion sickness medicine, cold medicine, antipyretics, analgesics, sedatives, antitussive expectorants, gastrointestinal analgesic antispasmodics, internal medicines containing antihistamine etc. (internal medicine for rhinitis, allergy medicine etc.)

- 2** Do not drive a motor vehicle or operate machinery after using this medicine. (Symptoms such as drowsiness, blurred vision, dazzling sunlight may occur.)

[Consult a doctor for the following]

- 1** The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
(2) A pregnant woman or a woman suspected of being pregnant
(3) Senior persons.
(4) A person with a medical history of being allergic to medicines and other substances.
(5) A person with the following symptoms:

Dysuria.

- (6) A person diagnosed as having the following:

Glaucoma, Heart disease.

- 3** The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese instructions written on this package case.

Related area	Symptoms
Skin	Rash · Redness, Itching
Central nervous system	Headache
Urinary system	Difficulty in urination
Others	Face hot flashes, Abnormal dazzling

- 4** The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and bring the Japanese instructions written on the package case, consult a doctor, pharmacist or registered distributor.

Thirst, Constipation, Sleepiness, Blurred vision.

Directions

Please take the following amounts and this medicine within 30 minutes before getting on board such as cars or ships for prevention of motion sickness.

Age	One dose	Number of doses per day
adult and children 15 years over	1 bottle(20mL)	once
children		Do not use

[When using this product]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children.
- (3) When taking additional doses as needed, take one dose at intervals of 4 hours or more. The number of daily doses is limited to 2 times.

Other information

- (1) Store in a cool and dry place free from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Please do not take products that have passed the expiration date.

Inactive ingredients: sucrose, citric acid, sodium benzoate, parabens and perfume

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00～12:00・13:00～17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.