

Product information

【Drug Facts】



<Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs **Medicine for motion sickness** (TABLETS)

KIDS TIMEL TABLETS

● "KIDS TIMEL TABLETS", taken once a day, reduce nervous excitement caused by vehicle sway and acceleration, and reduce or relieve the occurrence of dizziness, nausea and headache caused by motion sickness. If you use about 30 minutes before getting on a car, ship, airplane etc, you will be relieved from unpleasant motion sickness, thereby making travel and excursion more fun.

<i>Active ingredients (in 2 tablets/daily dose)</i>	<i>Purpose</i>
Meclizine HCl 33.3mg	Antiemetic
Scopolamine Hydrobromide Hydrate 0.166mg	Sedative
Anhydrous Caffeine 15mg	Headache treatment

Uses For the prevention and treatment of dizziness, nausea, headache associated with motion sickness.

Warnings Do not use for child under 3 years old.

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

1 While taking this medicine, please do not take any of the following medicines.

Other medicines such as motion sickness medicine, cold medicine, antipyretics, analgesics, sedatives, antitussive expectorants, gastrointestinal analgesic antispasmodics, internal medicines containing antihistamine etc. (internal medicine for rhinitis, allergy medicine etc.).

2 Do not drive a motor vehicle or operate machinery after using this medicine. (Symptoms such as drowsiness, blurred vision, dazzling sunlight may occur)

[Consultation]

1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A pregnant woman or a woman suspected of being pregnant.
- (3) Senior persons.
- (4) A person with a medical history of being allergic to medicines and other substances.
- (5) A person with the following symptoms:
Dysuria.
- (6) A person diagnosed as having the following:
Glaucoma, Heart disease.

2 The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Skin	Rash · Redness, Itching
Central nervous system	Headache
Urinary system	Difficulty in urination
Others	Face hot flashes, Abnormal dazzling

3 The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Thirst, Constipation, Sleepiness, Blurred vision

Directions

Please take the following amounts.

Age	One dose	Number of doses per day
children 11 years to under 15 years	2 tablets	once


children 3 years to under 11 years	1 tablet	(Please take this medicine within 30 minutes before getting on board such as cars or ships for prevention of motion sickness.)
infants under 3 years	Do not use	

[When using this product.]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) If you want to use this medicine to children over 3 years old, please be careful not to stuck it in their throat.
- (4) When using this medicine for the first time,

Please push the projection of the PTP sheet containing tablets strongly with your fingertips as shown below, break the aluminum sheet on the back side, push out and take it.

(If you mistakenly swallow the PTP sheet as it is, it leads to an unexpected accident such as stabbing into the esophagus mucosa)



- (5) This medicine can be drunk without water, but if you melt in the mouth, bitterness comes out, please swallow early.

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] **<Kyowa-Gifu-Japan>**

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.