

## Product information

**【Drug Facts】**



〈Kyowa-Gifu-Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs

**Antitussive expectorant**

(CAPSULES)

### NOSUPOL CHINGAI CAPSULE

“NOSUPOL CHINGAI CAPSULE” is a capsule that contains Dihydrocodeine phosphate, dl-Methyl ephedrine hydrochloride, Noscapine, and an antitussive component of guaifenesin. It will improve the symptoms of painful cough and sputum.

| <i>Active ingredient (in 6capsules/daily dose)</i> | <i>Purpose</i>                        |
|--|---------------------------------------|
| Dihydrocodeine phosphate 18.0mg·····               | Anti-cough                            |
| dl-Methylephedrine hydrochloride 40.0mg··          | Bronchodilator                        |
| Noscapine 60.0mg··········                         | Cough suppressant                     |
| Chlorpheniramine maleate 7.5mg······               | Reduce sneezing, runny or stuffy nose |
| Guaifenesin 300.0mg··········                      | Antitussive expectorant               |
| Anhydrous Caffeine 150.0mg········                 | Headache treatment                    |

#### **Uses**

Alleviation of the following respiratory function symptom:  
Cough and Sputum

#### **Warnings**

Do not use for children under 12 years old.

#### **[To be avoided]**

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

**1 While taking this medicine, please do not take any of the following medicines.**

Other antitussive expectorants, cold medicines, sedatives, oral medicines containing antihistamines (such as internal medicine for rhinitis, motion sickness medicine, medicine for allergy etc.)

- 2 Do not drive a motor vehicle or operate machinery after using this medicine. (Symptoms such as drowsiness, blurred vision, dazzling sunlight may occur)
- 3 You should not take this medicine during breastfeeding a baby. If you want take this medicine you should avoid to breastfeed your baby.
- 4 Do not take overdose or long-term continuously.

***[Consultation]***

1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A pregnant woman or a woman suspected of being pregnant
- (3) Senior persons.
- (4) A person who have ever had allergic symptoms due to any medication.
- (5) A person with the following symptoms:

High fever、 Dysuria

(6) A person diagnosed as having the following:

Heart disease, Hypertension, Diabetes, Glaucoma, Thyroid dysfunction, Respiratory dysfunction, Obstructive sleep apnea syndrome, Obesity

2 The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, stop the take immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

| Related area           | Symptoms                            |
|------------------------|-------------------------------------|
| Skin                   | Rash · Redness、 Itching             |
| Digestive organ        | Nausea · Vomiting, Loss of Appetite |
| Central nervous system | Dizzy                               |
| Urinary system         | Dysuria                             |

In very rare cases the following serious symptoms may occur. In this case, please get medical attention immediately.

| Symptoms' name | Symptoms |
|----------------|----------|
|----------------|----------|

|                        |   |
|------------------------|---|
| Aplastic anemia        | Bruising, nasal bleeding, Gum bleeding, Fever, Skin and mucous membranes appear pale, Fatigue, Palpitations, Shortness of breath, Mood worsening, Blood flowing, etc. |
| Agranulocytosis        | Sudden high fever, Chills, Sore throat etc.   |
| Respiratory depression | Shortness of breath, Breathlessness etc.  |

- 3** The following symptoms may appear after taking it, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.

Constipation, Thirst and Sleepiness.

- 4** If these symptoms do not improve after having used this medicine 5 or 6 times, stop taking the medicine, and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

### ***Directions***

Please take the following amount with water or lukewarm water after meals.


| Age                                 | One dose   | Number of doses per day |
|-------------------------------------|------------|-------------------------|
| adult and children 15 years over    | 2 capsules | 3 times                 |
| children 12 years to under 15 years | 1 capsule  |                         |
| children under 12 years             | Do not use |                         |

#### ***[When using this product]***

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) How to take out of capsules

Please push the projection of the PTP sheet containing capsules strongly with your fingertips as shown below, break the aluminum sheet on the back side, push out and take it.

(If you mistakenly swallow the PTP sheet as it is, it leads to an unexpected accident such as stabbing into the esophagus mucosa)



### ***Other information***

- (1) Store in a cool and dry with less humidity place free from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) Please do not take products that have passed the expiration date.

### ***Inactive ingredients:***

Potato starch, Lactose, Cellulose, Carmellose Ca, Hypromellose, FD&C Red No. 2[Red2G FCF], FD&C Red No. 102[New Coccine], And In the capsule body., Gelatin, FD&C Blue No.1[Brilliant Blue FCF], Red No. 102[New Coccine], Na lauryl sulfate, Na bisulfite

### ***Questions or comments***

If you have any suggestions about this product, please contact the shop you purchased or the following address.

---

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

### ***[Request of limited correspondence] <Kyowa-Gifu-Japan>***

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

### ***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan..
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.

- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.