Revision Sep 2020 No. 014

Product information

(Drug Facts)



Ҟ 〈Kyowa−Gifu−Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs

Athlete's foot • Ringworm

(LIQUID)

PALGRAN GX EKI

The athlete's foot and Ringworm are skin diseases caused by the fungus Trichophyton, which infests the stratum corneum of the skin, etc., often accompanied by itching and pain. "PALGRAN GX EKI" contains eight active ingredients. Once a day by using, it improves itching and painful (of athlete's foot and Ringworm) unpleasant symptoms.

Active ingredient (in 100mL)	Purpose
Butenafine hydrochloride 1.0g·····	Anti-trichophyton agent
Lidocaine 2.0g·····	Reduce itching with local anesthesia
Crotamiton 5.0g·····	Reduce itching
Glycyrrhetinic acid 0.5g·····	Reduce inflammation
dl-Camphor 1.0g·····	Give a refreshing feeling
Isopropyl methyl phenol 0.3g·····	Bactericidal action

Chlorpheniramine maleate 0.5g Reduce the occurrence of itching I-Menthol 1.0g · · · · · · · · · Give a refreshing feeling

[PRECAUTIONS CONCERNING INGREDIENTS]

Since this medicine contains alcohol, it may feel like sting at the time of application.

Uses

Athlete's foot (Tinea pedis), Ringworm in the groin, Ringworm in the body (Tinea corporis)

[Treatment points for Athlete's foot]

- Always clean the affected area. Let's use clean things such as socks and shoes,etc.
- Apply the medicine widely, because the infestation of Trichophyton often spreads from the range where the symptom appears

	because it may recur, so let's continue the	
treatment patiently for a while.		

O Let's prevent wetting of the affected area and try to keep it dry

Warnings

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions occurring.)

1. Do not use to the following person:

If you have had allergic symptoms (rash, redness, itching, edema etc.) due to this medicine or the components in it.

- 2. Do not use for the next part:
- (1) Around eyes or eyes, mucous membranes (oral cavity, nasal cavity, vagina, etc.), scrotum, vulva etc.
- (2) Eczema.
- (3) Severely affected areas of dampness, dripping, cracks and external wounds.

[Consultation]

- 1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.
- (1) A person undergoing medical treatment from a doctor.
- (2) A pregnant woman or a woman suspected of being pregnant.
- (3) Infants
- (4) A person who have ever had allergic symptoms due to any medication.
- (5) The affected area is on the face or in a wide area.
- (6) A person whose affected area is purulent.
- (7) A person who does not know clear whether "Eczema" or "Athlete's foot (Tinea pedis), Ringworm in the groin, Ringworm in the body (Tinea corporis) by trichomonas". (If you have symptoms such as itching or sore in the scrotum, there may be other causes than eczema etc.)
- The following symptoms may be adverse reactions to this medicine. If any
 of these symptoms occur, discontinue use immediately and consult a
 doctor, pharmacist, or registered distributor with the Japanese package
 insert.

Related area	Symptoms
Skin	Rash • redness, itching, blotch, swelling, irritation, heat on skin, skin scales, sores, dryness, irritability, crack

3. If symptoms do not improve after using it for about 2 weeks, stop using the medicine, bring this Japanese package insert, and consult a doctor, pharmacist or registered distributor.

Directions

Apply an appropriate amount to the affected area once a day

PRECAUTIONS CONCERNING DOSAGE AND ADMINISTRATION>


- (1) Please follow the recommended dosage and directions.
- (2) Please do not use while the affected area or its surroundings are dirty
- (3) Be careful not to get into eyes. If it gets in your eyes, please wash immediately with water or lukewarm water. And If the symptoms are severe, please be given medical treatment by an eye doctor immediately.
- (4) Do not touch your eyes or mucous membranes with this medicine.
- (5) If you want to use it for children, please use under the supervision of parents
- (6) Please use only for external treatment

[How to use the container]

 Before use, with the tip of the container facing up, press the tip of the plug with your finger two to three times to drain the air.

(When the product is placed in a hot place, when the temperature is high, or when the amount of fluid decreases, the internal pressure in container may increase, and too much chemical solution may be released.)

When using in the affected area, apply the tip gently to the affected area with the container facing downward or obliquely. If you remove the tip of container from the affected area, the medicine will not come out



(If you press strong or hold the container body firmly, a lot of medicine will be released.)

Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) Do not bring close to fire or put in fire. Also, do not throw used containers into fire.
- (5) Some agents in this medicine may soften synthetic resin (such as styrene) or dissolve paint, so please do not adhere on the floor or furniture etc.
- (6) Even within the expiration date after opening, please use as soon as possible from the point of quality maintenance.

Inactive ingredients:

Ethanol, 1,3-Butylene glycol, Macrogol, Phosphoric acid

Questions or comments

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Reception time 9:00~12:00·13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

(1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan..

- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the Revised date, the attached document is revised from time to time, so the English translation may not be the latest.