

## Product information

**【Drug Facts】**



＜Kyowa-Gifu-Japan＞

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 3 drugs      **ADHESIVE BANDAGE**      ( LIQUID TYPE )

### **EKITAI BANSOUKOU EX**

**【特徴】**

- 1 This medicine can quickly treat for "cracks", "akagire(cleft of skin)", "reverse stripping", "small cuts" and "scratches".
- 2 A thin film prevents from invading of bacteria, dust, etc. and from feeling of pain by getting wet.
- 3 Inconspicuous in transparent type.

**【Active ingredient】**

(in 100g)

**Effect**

Piroxirin 12g・

.....

Disinfection and protection for injury

**Inactive ingredients:**

Contains dl-camphor, castor oil, isopropyl myristate, ethyl acetate, and ethanol.

**【Indication】**

"cracks", "akagire(cleft of skin)", "reverse stripping", "small cuts" and "scratches".



### **Precautions**



**[When not to use the product]**

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

1. **Please** the medicine should not be used on the following parts:

(1) The eyes and periorbital area、mucosa (for example, oral cavity, nasal cavity or vagina, etc.)

- (2) Erosion, the purulent lesion
- (3) Large sores, deep wounds, eczema and bleeding areas.
- (4) Face, head.
- (5) Skin damaged by shaving, depilation, hair removal, bleaching, etc.



**[Consult a doctor for the following]**

**1. The following persons should consult a doctor, pharmacist, or registered salesperson before using this medicine.**

- (1) Patients undergoing medical treatment from a doctor.
- (3) Patients who have experienced allergic symptoms associated with drugs, etc.

**2. If the following symptoms appear after use, it may be a side effect, so stop using it immediately and consult your doctor, pharmacist, or registered salesperson with the Japanese instructions written on this package case.**

Related area	Symptoms
Skin	Rash/redness, itching, rash

**3. If symptoms do not improve after using between 5 and 6 days, stop using the medicine, consult a doctor, pharmacist, or registered salesperson registered seller with this case or the Japanese instructions written on this package case, and consult a doctor, pharmacist or registered salesperson.**

**【Directions】**

Clean the affected area, apply an appropriate amount only to the wound, and leave to dry.

**[When using this product]**

- (1) Do not use if the affected area or surrounding area is dirty.
- (2) If the affected area is wet with sweat or water, wipe off the moisture thoroughly before use.
- (3) Be careful not to get it in your eyes. If it does get in your eyes, immediately rinse with water or lukewarm water and consult an ophthalmologist.

- (4) This medicine should be given to children only under adult supervision.
- (5) This medicine should only be used externally and should never be taken orally.
- (6) Do not apply widely to areas other than the wound. (This may inhibit skin respiration.)
- (7) When you apply it to a wound, it will sting for a moment, but it will stop stinging once it dries.

### **【Precautions for storage and handling】**

- (1) Keep tightly sealed and store in a cool, low humidity place away from direct sunlight.  
  
(If the container is not sealed tightly, the medicine will harden and become unusable.)
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Keep away from fire.
- (5) Avoid use if there is a risk of accidental ingestion (e.g. by sucking finger).
- (6) Please be careful when trying to forcefully peel off the hardened film because you may injure your skin.  
  
\* If you need to remove this medicine quickly, apply layer the "EKITAI BANSOUKOU EX" on top and wipe off before it dries.
- (7) If this medicine gets on clothing, furniture, floors, etc., it is very difficult to remove and may cause discoloration, so please be careful when using it.
- (8) Do not use the product after the expiration date indicated on the outer box.

### ***Questions or comments?***

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

***[Request of limited correspondence]*** <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.