

Product information

【Drug Facts】



＜Kyowa-Gifu-Japan＞

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Transdermal Analgesic Anti-inflammatory (GEL)

VIPERCRON FB COOL GEL

- VIPERCRON FB COOL GEL is a non-steroid-based analgesic and anti-inflammatory medicine containing 3.0% of felbinac that acts directly on pain and swelling.
- Felbinac suppresses the formation of substances related to pain and swelling (prostaglandins) in the affected area and exerts an excellent effect on pain in joints, muscles, hips and shoulders.
- l-Menthol 6.0% formulation gives a pleasant refreshing sensation after application.

Active ingredients (in 100g)

Purpose

Felbinac 3.0g.....	Topical pain medication
l-Menthol 6.0g.....	To cool, Anti-inflammatory
Chlorpheniramine maleate 0.1g***	Anti-itching, Anti-inflammatory

Uses

Shoulder pain associated with stiff shoulders, Backaches, Joint pain, Muscle aches, Bruises, Sprains, Myelitis (pain or swelling at hand, wrist and ankle), Elbow pain (tennis elbow, etc.)

Warnings

Do not use for infants under 15 years old.

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions occurring.)

1. Do not use to the following person:

- (1) A person with a medical history of being allergic to this medicine or substances in it.
- (2) A person who has experienced asthma.

(3) A pregnant woman or a woman suspected of being pregnant.

(4) A child under 15 years old.

2. Do not use for the next part:

(1) Around eyes, mucous membranes etc.

(2) Eczema, rash, wound.

(3) The affected area is purulent on.

[Consult a doctor for the following]

1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

(1) A person undergoing medical treatment from a doctor.

(2) A person with a medical history of being allergic to medicines and other substances.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Skin	Rash • redness, Itching, Swelling, Irritation, Irritability

In rare cases, the following serious symptoms may occur. In that case, please get medical attention immediately.

Symptom name	Symptom
Shock(Anaphylaxis)	Immediately after using this medicine, itching of the skin, urticarial, hoarseness of voice, sneezing, throat itching, breathing, palpitations, turbulence of consciousness etc. appears.

3. If symptoms do not improve after using it about one week, stop using the medicine, bring the Japanese package insert, and consult a doctor, pharmacist or registered distributor.

Directions

Apply an appropriate amount to the affected area 2 to 4 times a day.

[When using this product]

(1) Do not use for children under 15 years old.

- (2) Please follow the recommended dosage and directions.
- (3) Be careful not to get into eyes. If it gets in your eyes, please wash immediately with water or lukewarm water. And If the symptoms are severe, please be given medical treatment by an eye doctor immediately.
- (4) Please use only for external treatment.
- (5) Please do not cover the affected area after using with a poorly breathable material (wrap film, straightening belt, etc.).

Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Do not bring close to fire or put in fire.
- (5) As it is volatile, please tighten the cap after use.
- (6) Be careful not to adhere to metal objects such as glasses, watches and accessories, clothing, plastics, leather products, and painted surfaces such as floors and furniture, as they may deteriorate or discolor.
- (7) Do not use products that have expired. Please use as soon as possible even within the expiration date after opening.

Inactive ingredients: Propylene glycol, Diisopropyl adipate, Sodium edetate hydrate, Hypromellose, Carboxyvinyl polymer, Diisopropanolamine and Ethanol.

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.

- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.