

# Product information

**【Drug Facts】**

 <Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

## **Cold Medicine (SYRUPS)**

### **NOSUPOL KODOMO KAZE SYRUP S**

"NOSUPOL KODOMO KAZE SYRUP S" is a cold medicine for children which contains Licorice extract and Bellflower flow extract from the crude drugs, in addition 5 kinds of active ingredients such as acetaminophen, dextromethorphan hydrobromide hydrate etc.

The function of the active ingredients relieves symptoms such as fever, sore throat, cough, phlegm, runny nose, stuffy nose, etc. occurred in child who tend to be exhausted by colds.

<i>Active ingredients (in 60 mL/Maximum daily dose)</i>	<i>Purpose</i>
---	----------------

Acetaminophen 300mg.....	Antipyretic analgesic
Chlorpheniramine maleate 2.4mg.....	Reduce sneezing, runny nose and stuffy nose
Dextromethorphan hydrobromide hydrate 16mg**	Reduce cough
dl-Methylephedrine hydrochloride 16mg.....	Bronchodilator
Anhydrous caffeine 32mg.....	Reduce headache
Licorice extract 50mg.....	Soothe sore throat and cough
Bellflower flower extract 0.24mg.....	Easier to expectorate phlegm

<b>Uses</b>	Relaxation of various symptoms of colds (runny nose, stuffy nose, sneeze, sore throat, cough, sputum, chills, fever, headache, joint pain, muscle pain)
-------------	---

<b>Warnings</b>	* In spite of this medicine for children, it describes the general precautions specified as a cold medicine.
-----------------	--

[To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

**1. Do not take to the following person:**

- (1) A person with a medical history of being allergic to this medicine or similar components in this medicine.
- (2) A person who have ever had asthma after taking this medicine or other cold medicine, antipyretic analgesics.

**2. While taking this medicine, please do not use any of the following medicines.**

Other medicines for cold, antipyretic analgesics, sedatives, antitussive expectorants, and internal medicines containing antihistamine etc. (internal medicine for rhinitis, motion sickness medicine, allergy medicine etc).

- 3. Do not drive the car or operate the machine after taking this medicine. (Symptoms such as sleepiness may occur)**
- 4. Do not drink alcohol before or after taking this medicine.**
- 5. Do not use in long-term continuously.**

***[Consult a doctor for the following]***

**1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.**

- (1) A person undergoing medical treatment from a doctor or dentist.
- (2) A pregnant woman or a woman suspected of being pregnant
- (3) A woman with breastfeeding.
- (4) A person who have ever had allergic symptoms due to any medication.
- (5) A person with the following symptoms:

High fever、 Dysuria

- (6) A person diagnosed as having the following:.

Thyroid dysfunction, diabetes, heart disease, hypertension, liver disease, kidney disease, stomach / duodenal ulcer, glaucoma.

**2 The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue take immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.**

Related area	Symptoms
Skin	Rash · Redness, itching
Digestive organ	Nausea · vomiting, loss of appetite
Central nervous system	Dizzy
Respiratory organ	Shortness, Stuffy of breath
Urology	Dysuria
Others	Excessive hypothermia

In very rare cases, the following serious symptoms may occur. In this case, please get medical attention immediately.

Related area	Symptoms
Shock(Anaphylaxis)	Immediately after taking medication, itching of the skin, urticarial, hoarseness of voice, sneezing, throat itching, breathing, palpitations, turbulence of consciousness etc. appears.
Skin mucosal eye syndrome (Stevens-Johnson syndrome), Toxic epidermal necrolysis, Acute pestiated pustule.	High fever, Redness of the eyes, Rashes, Sores on the lips, Sores in the throat, Extensive rash· redness on the skin, Small bumps (small blisters) on reddened skin. Heavy of Whole body. Persistent loss of appetite or Rapid worsening of symptoms.
Liver dysfunction	It shows fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, whole body dullness, loss of appetite, etc. appear.
Renal disorder	Fever, rash, decreased urine output volume, general whole body swelling or sagging, joint pain (nodal pain), diarrhea, etc. may appear.
Interstitial pneumonia	If you go up the stairs or you will run out of breath. And you may feel stuffy, empty cough, fever, etc. These will be suddenly appear or persist.
Asthma	When you breathe, the bronchus will squeeze with wheezing, whistling and a feeling of stuffiness appears.

Aplastic anemia	Bruising, nosebleeds, gum bleeding, fever, skin and mucous membranes appear pale, fatigue, palpitations, shortness of breath, mood worsening, bloodsheds etc. appear.
Agranulocytosis	Sudden high fever, chills, sore throat etc. appear.

**3 The following symptoms may appear after taking this medicine, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.**

Mouth thirst or Sleepiness.

**4 If these symptoms do not improve after having used this medicine 5 or 6 days, stop taking the medicine, and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.**

### ***Directions***

Please take the following amount with the attached scale with measuring cup, 3 times a day after meals and before occupation if necessary. In some cases, you can take up to 6 times a day at an interval of about 4 hours.

Age	One dose	Number of doses per day
3~7 years old	10mL	3 to 6 times
1~3 years old	7mL	
6 months more ~ under 1 year old	6mL	
3 ~6 months	5mL	
Less than 3 months	Do not use	

### ***[When using this product]***

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) To use for infants younger than 2 years, please priority to letting the doctor' consult, and so you will give this medicine only when you are unavoidable case.

### ***Other information***

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.

- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) Please wipe the mouth of the bottle after taking it, tighten the lid. If it is too bad, internal fluid may flow out or contaminate.
- (5) Please wash the measuring cup well after taking this medicine and keep it clean.
- (6) After opening once, please use up as early as possible from the point of quality preservation.
- (7) Please do not take medicines that have passed the expiration date

***[Request for storage]***

When the cap is tight, warm the cap with hot water (50–60 ° C), taking care not to enter hot water into the bottle. In that case, if the temperature of the hot water is too high, please be careful not to get burned and not to cause injury by broken the bottle.

***Inactive ingredients:***

Sugar, D – sorbitol, Na Benzoate, Paraben, Ethanol, Polyoxyethylene hardened castor oil, Caramel, Perfume, Glycerin, Vanillin, Propylene glycol.

***Questions or comments?***

If you have any suggestions about this product, please contact the shop you purchased or the following address.

---

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.

Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

***[Request of limited correspondence]***

**〈Kyowa-Gifu-Japan〉**

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan..

- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.