

Product information

【Drug Facts】



〈Kyowa-Gifu-Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs

Antitussive expectorant

(GRANULES)

NOSUPOL CHINGAI KARYU

”NOSUPOL CHINGAI KARYU” is one of granules wich have an effective combination of ingredients that suppress coughs and help cough up phlegm, and this granular formulation is dissolved quickly and is easy to take.

Active ingredient

Daily dose (3 packets) of this medicine

1g in one packet

Ingredient	Amount	Effect
Dihydrocodeine phosphate	30mg	Acts on the central nervous system to suppress coughs.
dl-methylephedrine hydrochloride	75mg	The effect of dilating the bronchi.
Chlorpheniramine maleate	7.5mg	The effect of against coughs caused by allergies.

Inactive ingredients

Contains Lactose hydrate, D-Mannitol, Potato starch, Saccharin Na hydrate, and Perfume .

Indication

Cough and Sputum.



Precautions



When not use the product

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

1 Do not take this medicine to the following person:

Children under 12 years old.

2 While taking this medicine, please do not take any of the following medicines.

Other expectorants, cold medicine, sedatives, or internal medicines containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, and medicine for allergies, etc.)

3 After taking this medicine, do not drive a car or operate machinery

(sleepiness may occur).

4 You should not take this medicine during breastfeeding a baby. If you want take this medicine you should avoid to breastfeed your baby.

Do not take this medicine in excess amount or for a long time.



Consult a doctor for the following

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

- (1) A person undergoing medical treatment from a doctor.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Senior persons.
- (4) A person who have ever had allergic symptoms due to any medication.
- (5) A person with the following symptoms:
High fever、 Dysuria.
- (6) A person diagnosed as having the following:.

Heart disease, hypertension, diabetes, glaucoma, thyroid dysfunction, respiratory dysfunction, obstructive sleep apnea syndrome, obesity.

2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this medicine, and show he Japanese package insert. to your physician, pharmacist, or registered salesperson for a consultation.

Related area	Symptoms
Skin	Rash · Redness、 Itching

Digestive organ	Nausea • Vomiting, Loss of Appetite
Central nervous system	Dizzy
Urinary system	Dysuria

In very rare cases, the following serious symptoms may occur. In this case, please get medical attention immediately.



Symptoms' name	Symptoms
Aplastic anemia	Bruising, nasal bleeding, Gum bleeding, Fever, Skin and mucous membranes appear pale, Fatigue, Palpitations, Shortness of breath, Mood worsening, Blood flowing, etc.
Agranulocytosis	Sudden high fever, Chills, Sore throat etc.
Respiratory depression	Shortness of breath, Breathlessness etc.

3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this medicine, and show the Japanese instructions written on this package case to your physician, pharmacist, or registered salesperson for a consultation.

Constipation, Dry mouth, Blurred vision

4. If these symptoms do not improve after having used this medicine 5 or 6 days, stop taking the medicine, and consult a doctor, pharmacist, or registered salesperson with the Japanese package insert.

Dosage and administration

Age	15 years over	12 years to under 15 years	under 12 years
One dose			Do not use
	1 packet	2/3 packet	

At a time up to three times a day, take this medicine avoid an empty stomach as much as possible, and in intervals of at least 4 hours

- When allowing children to take the medicine, a guardian must be present to watch and tell them how to take it.
- Please follow the prescribed usage and dosage.
- Do not give to children under 12 years of age.

Precautions for storage and handling

- (1) Store tightly closed and upright in a cool, dry place away from direct sunlight.
- (2) Keep this medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) Please do not take products that have passed the expiration date
- (5) If you want to divide one packet or take the rest, store the bag with the opening folded back and take it within 2 days.

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] **〈Kyowa-Gifu-Japan〉**

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.

- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.