

## Product information

**【Drug Facts】**



〈Kyowa-Gifu-Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it. (Revised in July 2022 due to changes in description guidelines)

Class ② drugs      **Antitussive expectorant**      (capsules)

### **NOSUPOL CHINGAI CAPSULE**

“NOSUPOL CHINGAI CAPSULE” is a capsule that contains Dihydrocodeine phosphate, dl-Methyl ephedrine hydrochloride, Noscapine, and an antitussive component of guaifenesin. It will improve the symptoms of painful cough and sputum.

#### **Active ingredient Daily dose (6 capsules) of this medicine**

| Ingredient                       | Amount  | Effect              |
|----------------------------------|---------|---------------------|
| Dihydrocodeine phosphate         | 18.0mg  | Antitussive effect  |
| dl-Methylephedrine hydrochloride | 40.0mg  | Bronchodilator      |
| Noscapine                        | 60.0mg  | Antiemetic effect   |
| Chlorpheniramine maleate         | 7.5mg   | Antihistamic effect |
| Guaifenesin                      | 300.0mg | Expectorant         |
| Anhydrous Caffeine               | 150.0mg | Headache treatment  |

#### **Inactive ingredients**

Contains Potato starch, Lactose, Cellulose, Carmellose Ca, Hypromellose, FD&C Red No. 2[Red2G FCF], FD&C Red No. 102[New Coccine], And In the capsule body., Gelatin, FD&C Blue No.1[Brilliant Blue FCF], Red No. 102[New Coccine], Na lauryl sulfate, Na bisulfite

#### **Indication**

Cough, Phlegm



#### **Precautions**

## When not use the product



(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

### 1 Do not take this medicine to the following person:

Children under 12 years old.

### 2 While taking this medicine, please do not take any of the following medicines.

Other expectorants, cold medicine, sedatives, or internal medicines containing antihistamines (internal medicine for rhinitis, medicine for motion sickness, and medicine for allergies, etc.)

### 3 After taking this medicine, do not drive a car or operate machinery

(Sleepiness may occur)

### 4 You should not take this medicine during breastfeeding a baby. If you want take this medicine you should avoid to breastfeed your baby.

### 5 Do not take this medicine in excess amount or for a long time.

## Consult a doctor for the following



### 1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

(1) A person undergoing medical treatment from a doctor.

(2) Pregnant women or women suspected of being pregnant.

(3) The elderly

(4) Patients who have experienced allergic symptoms associated with drugs, etc.

(5) A person with the following symptoms:

High fever、 Dysuria

(6) A person diagnosed as having the following:.

Heart disease, Hypertension, Diabetes, Glaucoma, Thyroid dysfunction, Respiratory dysfunction, Obstructive sleep apnea syndrome, Obesity

- 2** If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this medicine, and show the Japanese package insert to your physician, pharmacist, or registered salesperson for a consultation.

| Related area           | Symptoms                            |
|------------------------|-------------------------------------|
| Skin                   | Rash · Redness, Itching             |
| Digestive organ        | Nausea · Vomiting, Loss of Appetite |
| Central nervous system | Dizzy                               |
| Urinary system         | Dysuria                             |

In very rare cases, the following serious symptoms may occur. In this case, please get medical attention immediately.

| Symptoms' name         | Symptoms  |
|------------------------|---|
| Aplastic anemia        | Bruising, nasal bleeding, Gum bleeding, Fever, Skin and mucous membranes appear pale, Fatigue, Palpitations, Shortness of breath, Mood worsening, Blood flowing, etc. |
| Agranulocytosis        | Sudden high fever, Chills, Sore throat etc.   |
| Respiratory depression | Shortness of breath, Breathlessness etc.  |

- 3.** The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this medicine, and show the Japanese instructions written on this package case to your physician, pharmacist, or registered salesperson for a consultation.

Constipation, Thirst and Sleepiness.

- 4.** If these symptoms do not improve after having used this medicine 5 or 6 days, stop taking the medicine, and consult a doctor, pharmacist, or registered salesperson with the Japanese package insert.

## Dosage and administration

Please take the capsule directly with water or lukewarm water.

| Age                                 | One dose   | Number of doses per day |
|-------------------------------------|------------|-------------------------|
| adult and children 15 years over    | 2 capsules | 3 times                 |
| children 12 years to under 15 years | 1 capsule  |                         |
| children under 12 years             | Do not use |                         |


### <When using this product>

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.

- (3) When using this medicine for the first time,

<Capsule removal diagram>

Please push the projection of the PTP sheet containing capsules strongly with your fingertips as shown on the right, break the aluminum sheet on the back side, push out and take it.



(If you mistakenly swallow the PTP sheet as it is, it leads to an unexpected accident such as stabbing into the esophagus mucosa)

## Precautions for storage and handling

- (1) Store tightly closed and upright in a cool, dry place away from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) Please do not take products that have passed the expiration date

### *Questions or comments?*

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

***[Request of limited correspondence]***      **<Kyowa-Gifu-Japan>**

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.