

Product information

【Drug Facts】



〈Kyowa-Gifu-Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs **Transdermal Analgesic Anti-inflammatory (GEL)**

TAIANTERON FB PREMIUM GEL

■ characteristic

- VIPERCRON FB GOLD GEL is a non-steroidal analgesic and anti-inflammatory drug containing 3.0% Felbinac, which acts directly on the cause of pain and swelling.
- Contains 3.0% felbinac and 6.0% l-menthol.

Active ingredient (in 100g)

Ingredient	Amount	Effect
Felbinac	3.0g	Suppresses inflammation and relieves pain
l-Menthol	6.0g	Relieves pain and gives a refreshing feeling
Chlorpheniramine maleate	0.1g	Suppresses itching and inflammation

Indication

Shoulder pain associated with stiff shoulders, muscle pain, joint pain, lower back pain, tendonitis (pain and swelling in the hands, wrists, and ankles), elbow pain (tennis elbow, etc.), bruises, and sprains.



Precations



When not use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

1. **Please the medicine should not be used on the following parts:**

- (1) A person with a medical history of being allergic to this medicine or substances in it.

- (2) A person who has experienced asthma.
- (3) A pregnant woman or a woman suspected of being pregnant.
- (4) A child under 15 years old.

2. Do not use for the next part:

- (1) Around eyes, mucous membranes etc.
- (2) Eczema, rash, sores.
- (3) The affected areas with athlete's foot, ringworm, etc. or suppuration on.

3. Please do not use this medicine for a long time

 **Consult a doctor for the following**

1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) A person with a medical history of being allergic to medicines and other substances.

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.

Related area	Symptoms
Skin	Rash ▪ redness, Itching, Swelling, Irritation, Irritability

In rare cases, the following serious symptoms may occur. In that case, please get medical attention immediately.

Symptom name	Symptom
Shock(Anaphylaxis)	Immediately after using this medicine, itching of the skin, urticarial, hoarseness of voice, sneezing, throat itching, breathing, palpitations, turbulence of consciousness etc. appears.

3. If symptoms do not improve after using between 5 and 6 days, stop using the medicine, consult a doctor, pharmacist, or registered salesperson registered seller with this case or this Japanese product manual. the Japanese instructions written on this package case, and consult a doctor, pharmacist or registered salesperson.

Dosage and administration

Apply an appropriate amount to the affected area 2 to 4 times a day.

〈When using this product〉

- (1) Please follow the prescribed usage.
- (2) Be careful not to get it in your eyes. If it gets into your eyes, wash it immediately with cold or lukewarm water. If the symptoms are severe, please consult an ophthalmologist.
- (3) Please use only for external treatment.
- (4) Please do not cover the affected area after using with a poorly breathable material such as wrap film.

Precautions for storage and handling

1. Avoid high temperatures and store in a cool, dry place out of direct sunlight, tightly closed and upright.
2. Keep all medicine out of the reach of children.
3. Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
4. Do not keep this medicine away from fire. Also, do not throw it into a fire after use.
5. As it is volatile, please tighten the cap after use.
6. Be careful not to get it on metal items such as glasses, watches, accessories, synthetic clothing, plastics, painted surfaces such as floors or furniture, as it may cause deterioration or discoloration.
7. Do not use this medicine past their expiration date. Also, please use this as soon as possible after opening it, even if it is within the expiration date.

【Internal capacity】 50g

No fires

Class II petroleum

Danger class III

Inactive ingredients: Propylene glycol, carboxyvinyl polymer,
hydroxypropyl cellulose, edetate Na,
diisopropanolamine, ethanol

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.
Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.