Product information

[Drug Facts]



For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Skin Disease Treatment (OINTMENT)

TESITONFECT PLUS NANKOU

- © "TESITONFECT PLUS NANKOU" is a medicine that contains antedrug steroids and 8 ingredients in this medicine effect against insect bites, rashes, eczema and blotches.
- What is antedrug steroid?

Prednisolone valerate ester acetate exhibits excellent anti-inflammatory activity in the affected area, and it has the property of being weakly degraded when absorbed into the body afterward.

Active ingredients (in 1g)	Purpose
Prednisolone valeric acid ester acetic acid ester	A 1' ' Cl
(Antedrug steroids PVA) 1.5mg···	Anti-inflammatory ingredient
Glycyrrhetinic acid 5mg·····	Anti-inflammatory ingredient
Diphenhydramine hydrochloride 20mg····	Anti-histamine action
Crotamiton 50mg·····	Anti-itching ingredient
Lidocaine 20mg·····	Anti-itching ingredient
Tocopheryl acetate 10mg······	Blood circulation promoting ingredients
Allantoin 2mg·····	Remedies for skin
Isopropyl methyl phenol 1mg·····	Bactericidal ingredient

Eczema, Dermatitis, Heat rash, Rash, Itching, Insect bite,

Warnings

Uses

To be avoided?

Hives.

(Failure to observe the following may result in the current symptoms worsening or adverse reactions occurring.)

- 1. Do not use for the next part:
- (1) A person who has the affected areas with chicken pox (water pox), Athlete's foot (Tinea pedis), Ringworm in the groin, etc. and/or who have purulent on the affected part.
- (2) Around eyes, mucous membranes etc.
- 2. Please do not use it extensively on the face.
- 3. Do not use long-term continuously.

[Consult a doctor for the following]

- 1. The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.
- (1) A person undergoing medical treatment from a doctor.
- (2) A pregnant woman or a woman suspected of being pregnant.
- (3) A person who have ever had allergic symptoms due to any medication.
- (4) The affected area is on the face or spreaded on the wide.
- (5) A person with wet or awful sore on the affected part.
- 2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese instructions written on this package case.

Related area	Symptoms
Skin	Rash • Redness, Itching, Swelling, Skin irritation, Skin dryness, Stimulating or Heating on Skin, Irritability.
Skin (Affected area)	Athlete's foot (Tinea pedis) - Ringworm in the groin by trichomonas, Acne, Purulent symptoms, persistent feeling of irritation

3. If symptoms do not improve after using between 5 and 6 days, stop using the medicine, bring the Japanese instructions written on this package case, and consult a doctor, pharmacist or registered distributor.

Directions

Apply an appropriate amount to the affected area several times a day.

[When using this product]

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) Be careful not to get into eyes. If it gets in your eyes, please wash immediately with water or lukewarm water. And If the symptoms are severe, please be given medical treatment by an eye doctor immediately.
- (4) Please use only for external treatment

Other information

- (1) Store in a cool and dry with less humidity place free from direct sunlight by a seal tightly.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Do not use the product that has passed its expiration date (listed on the outer box).

Inactive ingredients:

Macrogol and Citric acid hydrate.

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Reception time 9:00~12:00·13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

(1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.

- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.