

## Product information

**【Drug Facts】**



〈Kyowa-Gifu-Japan〉

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs **Medicine for motion sickness (TABLETS)**

### TIMEL TABLETS

Timel Tablets, taken once a day, calms the nervous excitement caused by vehicle motion and acceleration, and suppresses or alleviates dizziness, nausea, and headaches caused by vehicles. If you take this medicine about 30 minutes before boarding a car, boat, airplane, etc., you will be relieved from unpleasant motion sickness and will enjoy your travels and hobbies even more.

#### **【Active ingredient】 Daily dose (1tablet) of this medicine**

Ingredient	Amount	Effect
Meclizine HCl	25mg	Antiemetic effect
Scopolamine Hydrobromide Hydrate	0.25mg	Sedative effect
Anhydrous Caffeine	40mg	Headache treatment

#### **【Inactive ingredients】**

Contains Saccharin Sodium, Lactose, Potato Starch, Calcium Stearate, Cellulose and FD&C Blue No.1[Brilliant Blue FCF] .

**【Indication】** For the prevention and treatment of dizziness, nausea, headache associated with motion sickness.

### **Precautions**

#### **When not use the product**

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

- 1 While taking this medicine, please do not take any of the following medicines.

Other medicines such as motion sickness medicine, cold medicine, antipyretics, analgesics, sedatives, antitussive expectorants, gastrointestinal analgesic antispasmodics, internal medicines containing antihistamine etc. (internal medicine for rhinitis, allergy medicine etc.)

## 2 After taking this medicine, do not drive a car or operate machinery

(symptoms such as sleepiness, blurred vision and abnormal brightness may occur).



### Consult a doctor for the following

#### 1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.

- (1) Patients undergoing medical treatment from a physician or dentist.
- (2) Pregnant women or women suspected of being pregnant.
- (3) The elderly
- (4) Patients who have experienced allergic symptoms associated with drugs, etc.
- (5) Persons with the following symptoms:

urination difficulty

#### (6) Persons diagnosed as having the following'

Glaucoma, heart disease

#### 2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this medicine, and show the Japanese package insert to your physician, pharmacist, or registered salesperson for a consultation.

Related area	Symptoms
Skin	Rash ▪ redness, Itching
Central nervous system	Headache
Urinary system	Difficulty in urination
Others	Face hot flashes, Abnormal dazzling

3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this medicine, and show the Japanese instructions written on this package case to your physician, pharmacist, or registered salesperson for a consultation.

Dry mouth, constipation, drowsiness, blurred vision

## 【Dosage and administration】

Please take the following amounts


Age	One dose	Number of doses per day
adult and children 15 years and over	1 tablet	once
children 7 years to under 15 years	1/2 tablet	(To prevent motion sickness, take 30 minutes before getting on a car or boat.)
children under 7 years	Do not use	

### 〈When using this product〉

- (1) Please follow the recommended dosage and directions.
- (2) This medicine should be given to children only under adult supervision.
- (3) When using this medicine for the first time,

Please push the projection of the PTP sheet containing tablets strongly with your fingertips as shown below, break the aluminum sheet on the back side, push out and take it.

(If you mistakenly swallow the PTP sheet as it is, it leads to an unexpected accident such as stabbing into the esophagus mucosa)



- (4) This medicine can be drunk without water, but if you melt in the mouth, bitterness comes out, please swallow early.

## **【Precautions for storage and handling】**

- (1) Store tightly closed and upright in a cool, dry place away from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) If you will use the rest half of one tablet, wrap it in the clean paper, store in the outer box, and take it within 2 days.
- (5) Please do not take products that have passed the expiration date

## ***Questions or comments?***

If you have any suggestions about this product, please contact the shop you purchased or the following address.

---

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

### ***[Request of limited correspondence]*** <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

## ***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.

- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.