

# Product information

## 【Drug Facts】



<Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

## Cold Medicine (FINE PARTICLES)

### SOUGOU KAZE KUSURI GOLD A <BIRYU>

“SOUGOU KAZE KUSURI GOLD A <BIRYU>”, seven kinds of active ingredients are effective for various symptoms of cold, such as headache, fever, cough, runny nose.

#### *Active ingredients (in 3 packets/daily dose) Purpose*

( 1.200mg in 1 Packet )

|  |  |
|--|--|
| Acetaminophen 900mg.....                 | Antipyretic analgesic                        |
| Dihydrocodeine phosphate 24mg .....      | Anti-cough                                   |
| Chlorpheniramine maleate 7.5mg.....      | Reduce sneezing, runny nose and stuffy nose. |
| dl-Methylephedrine hydrochloride 60mg .. | Bronchodilator                               |
| Anhydrous Caffeine 75mg.....             | Headache treatment                           |
| Guaifenesin 180mg.....                   | Antitussive expectorant                      |
| Riboflavin 12mg .....                    | Vitamin B2 supplementation                   |

#### *Uses*

Alleviation of cold symptoms such as runny nose, stuffy nose, sneezing, sore throat, cough, sputum, chills (since due to fever), fever, headache, joint pain, muscle pain.

#### *Warnings*

Do not use for children under 12 years old.

#### [To be avoided]

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

#### 1. Do not take to the following person:

- (1) A person with a medical history of being allergic to this medicine or substance in it.

(2) A person who have ever had asthma taking this medicine or other cold medicine, antipyretic analgesics.

(3) Children under 12 years old.

**2 While taking this medicine, you should not use in combination with the following medicines:**

Other medicines for cold, antipyretic analgesics, sedatives, antitussive expectorants, and internal medicines containing antihistamine etc. (internal medicine for rhinitis, motion sickness medicine, allergy medicine etc).

**3. Do not drive a motor vehicle or operate machinery after using this medicine. (Symptoms as drowsiness may occur)**

**4. You should not take this medicine during breastfeeding a baby. If you want take this medicine you should avoid to breastfeed your baby.**

**5. Do not drink alcohol before or after taking this medicine.**

**6. Do not use long-term continuously.**

***[Consult a doctor for the following]***

**1 The following persons should consult a doctor, pharmacist, or registered distributor before using this medicine.**

(1) A person undergoing medical treatment from a doctor or dentist..

(2) A pregnant woman or a woman suspected of being pregnant.

(3) Senior persons.

(4) A person with a medical history of being allergic to medicines and other substances.

(5) A person with the following symptoms:

High fever, Dysuria.

(6) A person diagnosed as having the following:

Thyroid dysfunction, diabetes, heart disease, hypertension, liver disease, glaucoma, systemic lupus erythematosus, mixed connective tissue disease, respiratory dysfunction, obstructive sleep apnea syndrome, obesity.

**3. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered distributor with the Japanese package insert.**

|              |          |
|--------------|----------|
| Related area | Symptoms |
|--------------|----------|

|                        |   |
|------------------------|---|
| Skin                   | Rash / redness, itching, blue bruise  |
| Digestive organ        | Nausea / vomiting, loss of appetite, stomach discomfort, stomach pain, stomatitis, heartburn, stomach upset, bleeding, abdominal pain, diarrhea, bloody stool |
| Central nervous system | Dizziness, excitement, convulsions  |
| Urology                | Difficulty urinating  |
| Others                 | Eye blur, tinnitus, swelling, nose bleeding, bleeding of gums, bleeding is difficult to stop, bleeding, back pain, excessive temperature drop, tired          |

In very rare cases the following serious symptoms may occur. In this case, please get medical attention immediately.

| Symptoms' name  | Symptoms   |
|---|--|
| Shock(Anaphylaxis)  | Immediately after taking medication, itching of the skin, urticaria, hoarseness of voice, sneezing, throat itching, breathing, palpitations, turbulence of consciousness etc. appears.   |
| Skin mucosal ocular syndrome(Stevens–Johnson syndrome) , Toxic epidermal necrolysis | High fever, redness of the eyes, sore lips, sore throat, sore throat, extensive skin rash / redness, small rashes on the reddish skin, general dullness, lack of appetite, etc. or these symptoms get worse rapidly.                       |
| Liver dysfunction   | It shows fever, itching, rash, jaundice (yellowing of the skin and whites of the eyes), brown urine, whole body dullness, loss of appetite, etc. appear.   |
| Renal disorder  | Fever, rash, decreased urine volume, whole body swelling, whole body sagging, joint pain (sore throat), diarrhea, etc. appear.   |
| Aseptic meningitis  | Severe headaches accompany with the neck strain, and fever, nausea, vomiting etc. may occur. (Such symptoms are often reported, especially in people who are treated for systemic lupus erythematosus or mixed connective tissue disease.) |
| Interstitial pneumonia  | If you go up the stairs or do a little impossible, you will be out of breath, Nausea / vomiting and feel stuffy, fever etc. These are suddenly appeared or persisted.  |
| Asthma  | When you breathe, the bronchus will squeeze with wheezing, whistling and a feeling of stuffiness appear.   |

|                        |   |
|------------------------|---|
| Aplastic anemia        | Bruising, nosebleeds, gum bleeding, fever, skin and mucous membranes appear pale, fatigue, palpitations, shortness of breath, mood worsening, bloodsheds etc. appear. |
| Agranulocytosis        | Sudden high fever, chilling, sore throat etc. appear.   |
| Respiratory depression | Constipation, Shortness of breath, breathing difficulties and so may appear.  |

4. The following symptoms such as Constipation, Thirst, Sleepiness may appear after taking this medicine, so if such symptoms persist or increase. Please stop taking it and consult a doctor, pharmacist or registered distributor with the Japanese package insert.
5. If these symptoms are not good even after taking between 5 and 6 times, please stop taking the medicine and consult your doctor, pharmacist or registered distributor with the Japanese package insert.

### ***Directions***

Please take 1 time as possible within 30 minutes after meals, on three times a day.

| Age   | One dose   | Number of doses per day |
|---|------------|-------------------------|
| adult and children 15 years old and over    | 1 packet   | 3 times after meal      |
| children 12 years old to under 15 years old | 2/3 packet |                         |
| children under 12 years old.                | Do not use |                         |

#### ***[When using this product]***

- Please follow the recommended dosage and directions.
- This medicine should be given to children only under adult supervision.
- If you take medicine after dividing one packet, please fold the mouth of the packet and store it, please take it within 2 days.

### ***Other information***

- (1) Store in a cool and dry with less humidity place free from direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)

(4) Do not take this medicine after the expiration date on the outer box.

***Inactive ingredients:***

Cellulose, Anhydrous silicic acid, Potato starch, D-Mannitol, Hydroxypropylcellulose, Magnesium aluminate metasilicate, Aspartame, Fragrance, Vanillin.

***Questions or comments?***

If you have any suggestions about this product, please contact the shop you purchased or the following address.

---

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.  
Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00•13:00~17:00 (Excluding weekends and holidays)

***[Request of limited correspondence]*** <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. ([www.kyowa-ci.co.jp](http://www.kyowa-ci.co.jp)).

***Disclaimer statement***

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.