

Product information

【Drug Facts】



〈Kyowa-Gifu-Japan〉

For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Class ② drugs **Skin Disease Treatment** (LIQUID)

TESITON HR LIQUID

Active ingredients

(in 100mL)

Effect

Prednisolone valeric acid ester acetic acid ester
(PVA) 0.15g····

Reduce skin irritation

Diphenhydramine hydrochloride 2.0g····

Reduce the occurrence of itching

l-Menthol 3.5g····

Give a refreshing feeling

l-Camphor 1.0g····

Give a refreshing feeling

Isopropyl methyl phenol 0.1g····

Bactericidal ingredient

Indication

Eczema, dermatitis, heat rash, rash, itching, insect bites, hives



Precautions



When not to use the product

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

1. Please the medicine should not be used on the following parts:

- (1) Varicella (chickenpox)、Athlete's foot/tinea or purulent lesions
- (2) Wound surface、Periorbital area、Mucosa

2. Please do not use extensively on the face

3. Please do not use this medicine for a long time



Consult a doctor for the following

1. The following persons should consult a doctor, pharmacist, or registered salesperson before using this medicine.

- (1) Patients undergoing medical treatment from a doctor.
- (2) Pregnant women or women suspected of being pregnant.
- (3) Patients who have experienced allergic symptoms associated with drugs, etc.
- (4) Those with extensive lesions.
- (5) Those with severe moistening or erosion

2. If the following symptoms appear after use, it may be a side effect, so stop using it immediately and consult your doctor, pharmacist, or registered salesperson with this case.

Related area	Symptoms
Skin	Rash ▪ redness, itching, swelling
Skin (Affected area)	Ringworm such as athlete's foot and ringworm, acne, suppurative symptoms, and persistent irritation

3. If symptoms do not improve after using between 5 and 6 days, stop using the medicine, consult a doctor, pharmacist, or registered salesperson registered seller with this case or this Japanese product manual. the Japanese instructions written on this package case, and consult a doctor, pharmacist or registered salesperson.

Dosage and administration

Apply an appropriate amount to the affected area several times a day.

When using this product

- (1) Remove the cap and lightly press the sponge part against your skin several times to soak the medicinal solution into the sponge before use. If the chemical solution does not soak into the sponge surface sufficiently, the sponge surface may be damaged.

After using this medicine, be sure to close the cap tightly to prevent the chemical solution from drying out on the sponge.

- (2) Please follow the prescribed usage and dosage.
- (3) This medicine should be given to children only under adult supervision.
- (4) Be careful not to get it in your eyes. If it gets into your eyes, immediately wash it with water or lukewarm water. If the symptoms are severe (such as persistent redness, pain, or unstopable tearing), please consult an ophthalmologist.
- (5) Please use only for external treatment.

Precautions for storage and handling

- (1) Store upright and tightly closed in a cool place out of direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)
- (4) Please keep away from fire. Also, do not throw the used amount into fire.
- (5) Be careful not to get it on glasses, watches, accessories, plastics, synthetic fiber products, leather products, painted surfaces such as floors or furniture, as it may cause deterioration.
- (6) Do not use the medicine past their expiration date. Also, please use the medicine product as soon as possible after opening it, even if it is within the expiration date.

【Capacity】 40mL

No fires
Class II petroleum, danger class III
(ethanol)

Inactive ingredients:

Contains dibutylhydroxytoluene, 1,3-butylene glycol, and enanol.

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.



Customer consultation room ☎ 058-246-8771

Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.