

Product information

【Drug Facts】**〈Kyowa-Gifu-Japan〉**

For resource saving the case is combined with the Japanese instruction written on the package case. Be sure to read the instruction carefully before use. Also, please keep it in a safe place so that you can read again, when you need it.

Class ② drugs

Antipruritic and Anti-inflammatory Drugs

(LIQUID)

TESITON EH LIQUID

When applying medicine to the skin inside the ear, do not insert the cotton swab deep into the ear, as this may damage the inner part of the ear or the eardrum.

Hold the cotton ball about 1.5cm from the tip (underneath the cotton ball) and apply gently without rubbing.

Precautions when using in ears


- (1) This medicine is not an ear drop, so do not drop it directly into the ear.
- (2) When using on the skin inside the ear, be sure to follow the instructions on the back of the container.
- (3) Please do not use it for a long period of time. (As a guideline, within 2 weeks)
- (4) If you suspect that your eardrum is perforated or have ear discharge, consult a doctor, pharmacist, or registered salesperson before use.
- (5) If you have difficulty hearing after using this medicine, stop using it and consult a doctor, pharmacist, or registered seller with this case or this Japanese product manual.
- (6) The recommended age to start using this medicine is 10 years or older. Please do not use this product on children (under 15 years old).

(7) Do not use earphones or hearing aids immediately after applying this medicine, and make sure they are completely dry before using. (After 1 hour as a guide)

Active ingredients (in 100mL)	Effect
Prednisolone valeric acid ester acetic acid ester (Antedrug steroids PVA) 0.15g***	Reduce skin irritation
Diphenhydramine hydrochloride 2.0g****	Reduce the occurrence of itching
Crotamiton 3.0g*****	Reduce itching
Dipotassium glycyrrhizinate 0.2g*****	Anti-inflammatory ingredient
l-Menthol 3.5g*****	Give a refreshing feeling
Lidocaine 1.0g*****	Reduce itching with local anesthesia
Isopropyl methyl phenol 0.1g*****	Bactericidal ingredient
Allantoin 0.2g*****	Remedies for skin

Indication Itching, eczema, dermatitis, rash, hives, heat rash, insect bites

 **Precautions**

 **When not to use the product**

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

1. Please the medicine should not be used on the following parts.

- (1) Varicella (chickenpox), Athlete's foot/tinea or purulent lesions
- (2) Wound surface, Periorbital area, Mucosa

2. Please do not use extensively on the face

3. Please do not use this medicine for a long time

 **Consult a doctor for the following**

1. The following persons should consult a doctor, pharmacist, or registered salesperson before using this medicine.

- (1) Patients undergoing medical treatment from a doctor.

- (2) Pregnant women or women suspected of being pregnant.
 - (3) Patients who have experienced allergic symptoms (Rash, redness, rash, etc.) associated with drugs, etc.
 - (4) Those with extensive lesions.
 - (5) Those with severe moistening or erosion
- 2. If the following symptoms appear after use, it may be a side effect, so stop using it immediately and consult your doctor, pharmacist, or registered salesperson with this case.**

Related area	Symptoms
Skin	Rash ▪ redness, itching, swelling, blotch, irritation, heat on skin, dryness, irritability
Skin (Affected area)	Ringworm such as athlete's foot and ringworm, acne, suppurative symptoms, and persistent irritation

- 3. If symptoms do not improve after using between 5 and 6 days, stop using the medicine, consult a doctor, pharmacist, or registered salesperson registered seller with this case or this Japanese product manual. the Japanese instructions written on this package case, and consult a doctor, pharmacist or registered salesperson.**

[Dosage and administration]

Apply an appropriate amount to the affected area several times a day.

When using this product

- (1) Please follow the prescribed usage and dosage.
- (2) This medicine should be given to children only under adult supervision.
- (3) Be careful not to get it in your eyes. If it gets into your eyes, immediately wash it with water or lukewarm water. If the symptoms are severe (such as persistent redness, pain, or unstoppable tearing), please consult an ophthalmologist.
- (4) This medicine is for external use only and should not be taken internally.

(5) If the affected area is scarred, you may feel a sting when you apply it.

How to use the container

(1) Before use, with the tip of the container facing upward, press the tip of the inner stopper 2 to 3 times with your fingers to release the air inside.

(This medicine is to prevent the pressure inside the container from increasing when the container is placed in a hot place, or when the chemical solution is running low, causing a large amount of the chemical solution to come out.)

(2) When using, hold the container top on facing down or diagonally and lightly press the tip onto the affected area to apply. When you remove the tip of the container from the affected area, the liquid will no longer come out.

(If you press it too hard or squeeze the container body too tightly, more liquid will come out.)

Precautions for storage and handling

(1) Store in a cool place free from direct sunlight by a seal tightly.

(2) Keep all medicine out of the reach of children.

(3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality.)

(4) Keep away from fire. Also, do not throw it into a fire after use.

(5) Be careful not to get it on metal objects such as glasses, watches, accessories, plastics, synthetic clothing, leather products, or painted surfaces, and be careful such as floors or furniture, so these cases may cause deterioration.

(6) If this medicine gets on dyed hair, the color may fade.

(7) Tighten the cap after use.

(8) Do not use this product after the expiration date (written on the bottom of the box).

Don't put it in your eyes

No fires
Class II petroleum, danger class III
(ethanol)

Inactive ingredients: Contains 1,3-butylene glycol, dibutylated hydroxytoluene, ethanol, PH regulator, and hydrophobized hydroxypropyl methyl callose.

Contact information for the Relief System for Side Effects

Pharmaceuticals and Medical Devices Agency

http://WWW.pmda.go.jp/kenkouhigai_cmp/index.html

Telephone: 0120-149-931 (toll-free)

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.

Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence] <Kyowa-Gifu-Japan>

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.

- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.