

Product information

【Drug Facts】



<Kyowa-Gifu-Japan>

Please be sure to read this product information carefully before use. Keep this product information in a safe place so that you can read again, when you need it.

Class 2 drugs **Athlete's Foot and Ringworm Remedies** (LIQUID)

PALGRAN S EKI

Pargulan S liquid contains bifonazole, which has a long-lasting bactericidal effect against ringworm, etc., and is effective when used once a day.

Furthermore, four types of ingredients (lidocaine, crotamiton, glycyrrhetin, and l-menthol) can suppress unbearable itching and inflammation.

It is excellent quick-drying liquid with a smooth and refreshing feeling..

Active ingredient (in 100mL)

Bifonazole	1.0g	Continuously inhibits the growth of tinea fungus.
Lidocaine	2.0g	Relieve itching and pain.
Crotamiton	5.0g	Relieve itching.
Glycyrrhetic acid	0.5g	Suppresses inflammation.
l-Menthol	2.0g	To reduce inflammation and give refreshing of the affected area.

【Indication】 Alleviation of the symptoms by Athlete's foot (Tinea pedis)、 Ringworm in the groin、 Ringworm in the body (Tinea corporis)

Points of athlete's foot treatment

1. Cleanliness

To suppress the growth of ringworm and prevent secondary infection, wash the affected area thoroughly and keep it clean by wiping their moisture to dry.

2. Apply over a wide area

Trichophyton, which is the cause of athlete's foot and ringworm, often spreads beyond the area where the symptoms appear, so make sure to apply the medication over a wide area.

3. Dryness

Choose socks and footwear that are breathable to prevent stuffiness and keep you dry.

4. Continuous treatment

Even if symptoms disappear such as itching, to continue the treatment for at least one season.



Precautions



When not use the product

(Failure to observe the following may result in the current symptoms worsening or adverse reactions/ accidents occurring.)

1. Do not use for the next part:

- (1) Eyes and the area around the eyes or mucous membranes (e.g., oral cavity, nasal cavity, vagina, etc.), scrotum, vulva, etc.
- (2) Eczema
- (3) Severely moist, sore, cracked or damaged areas.



Consult a doctor for the following

1 The following persons should consult a doctor, pharmacist, or registered salesperson before using this medicine.

- (1) A person undergoing medical treatment from a doctor.
- (2) Infants
- (3) A person with a medical history of being allergic to medicines and other substances.
- (4) The affected area is on the face or in a wide area.
- (5) The affected area is purulent on.
- (6) A person who are unsure whether they have eczema or athlete's foot, jock itch, or jock itch.

(If you have symptoms such as itching or sores on your scrotum, it is often due to other causes such as eczema.)

2. The following symptoms may be adverse reactions to this medicine. If any of these symptoms occur, discontinue use immediately and consult a doctor, pharmacist, or registered salesperson with the Japanese package insert.

Related area	Symptoms
Skin	Rash ▪ redness, itching, blotch, swelling, irritation, heat on skin, dryness, irritability

3. If symptoms do not improve after using it for about 2 weeks, stop using the medicine, bring this Japanese package insert, and consult a doctor, pharmacist or registered salesperson.

【Dosage and administration】

Clean the affected area and apply an appropriate amount to the affected area once a day


《PRECAUTIONS CONCERNING DOSAGE AND ADMINISTRATION》

- (1) Please do not use while the affected area or its surroundings are dirty
- (2) Be careful not to get it in your eyes. If it comes into contact with your eyes, immediately rinse with cold or lukewarm water and seek medical attention from an ophthalmologist immediately.
- (3) If you want to use it for children, please use under the supervision of parents
- (4) Please use only for external treatment
- (5) Please strictly follow the prescribed usage and dosage.

How to use

- ① Before use, point the tip of the container upwards, press it with your finger and pull out the air inside

(This is to prevent internal pressure from rising and too much medical solution coming out when the medicine is placed in a high temperature place during hot weather or when the medical solution is low.)



- ② When using in the affected area, apply the tip gently to the affected area with the container facing downward or obliquely.



(If you remove the tip from the affected area, the medicinal solution will no longer come out.)

【Precautions for storage and handling】

- (1) Avoid high temperatures and store tightly closed in a cool place out of direct sunlight.
- (2) Keep all medicine out of the reach of children.
- (3) Do not transfer this medicine to a different container. (In this case, cause misuse or change quality)
- (4) Please do not get close to fire.
- (5) Please do not touch your eyes or mucous membranes with your hand wearing this medicine.
- (6) Do not use this medicine past their expiration date. Also, please use the product as soon as possible after opening it, even if it is within the expiration date.
- (7) Avoid getting this medicine on metal objects such as glasses, watches, and accessories, and on objects that may be deteriorated or discolored by alcohol or oils (plastic products, chemical fibers, leather, furniture, floors, etc.).

【Internal capacity】 30mL

Inactive ingredients: Sodium hyaluronate, urea, propylene glycol, diisopropyl adipate, lauromacrogol, ethanol

Contact information for the Relief System for Side Effects

Pharmaceuticals and Medical Devices Agency

http://WWW.pmda.go.jp/kenkouhigai_cmp/index.html

Telephone: 0120-149-931 (toll-free)

Questions or comments?

If you have any suggestions about this product, please contact the shop you purchased or the following address.

Consumer consultation window: Kyowa Pharmaceutical Industry Co., Ltd.

Customer consultation room ☎ 058-246-8771



Reception time 9:00~12:00・13:00~17:00 (Excluding weekends and holidays)

[Request of limited correspondence]

〈Kyowa-Gifu-Japan〉

- (1) In case of telephone consultation, please only in Japanese for a while.
- (2) The latest Japanese package insert is available on the website of Kyowa Pharmaceutical Co., Ltd. (www.kyowa-ci.co.jp).

Disclaimer statement

- (1) We use a format similar to the FDA OTC Drug Regulations, but this is not an OTC Drug approved under the FDA. Please note that it is licensed for domestic use in accordance with the Pharmaceutical Affairs Law of Japan.
- (2) This English translation was created by a manufacturer and distributor, but Japanese is a correct sentence and it should be as a reference material. In the case of any discrepancy between the two versions, the original Japanese version prevails.
- (3) The provider of this translation shall not be responsible for any damage that may be attributed to the use of the translation.
- (4) Do not use for commercial purpose reproduction, publication etc. of this English translation.
- (5) Although the English translation is prepared on the basis of the Japanese version of the attached document on the year and month, the attached document is revised from time to time, so the English translation may not be the latest.